American Heart Association/American Stroke Association
Strategic Policy Agenda
2014-2017

I. Introduction

The American Heart Associationôs public policy agenda provides our federal, state and local advocacy staff with strategic guidance and direction on policy issues and positions that align with and support the Associationôs mission and strategic priorities. The document attempts to capture the breadth of the Associationôs policy portfolio over the next three years, however, it is possible that issues will emerge during that time that could not be forecasted and might become a priority for the American Heart Association and incorporated into our work. Additionally, the Association scans the political landscape annually to identify leading opportunities and establishes federal and state priorities that serve to focus our immediate advocacy efforts on those issues that present the greatest opportunity for success in achieving mission and strategic priority

ationôs commitment to eliminate heaea

The American Heart Associationôs Advocacy Coordinating Committee (AdCC), a committee of the Associationôs national board, is responsible for establishing the Associationôs policy positions, public policy agenda, and annual legislative and regulatory priorities. The public policy agenda and annual priorities are a product of a rigorous internal process that is informed by our science, guided by our 2020 health impact goal and strategic plan, and refined through the advice and counsel provided by AHA staff and volunteers. These priorities, which are

The associationôs research priority includes all forms of scientific studies, including basic science as well as clinical, translational, health services (outcomes), genomics, and comparative effectiveness research and the overall research environment. Effectively preventing and treating disease depends on accurate knowledge about its causes, on how disease affects the body, on drugs that combat disease, on devices that are safe and work, and on operations that cure as well as clinical research that helps enable health care professionals to assist their patients and their families in building the skills they need to adopt and maintain a healthy lifestyle. The knowledge, material and skills on which prevention and treatment are based have come from a variety of sources, including information that can only be obtained from research on both animals and humans. Animal research has improved the health and welfare of both animals and humans. The decline in death rates in the United States from heart disease and stroke since the 1960s is due to lifestyle changes and new methods of treatment and prevention, many of which are based on animal research. The association generally opposes legislation and regulations that would curtail necessary heart disease and stroke research or make it unduly difficult or costly.

Demographics

Death rates from coronary heart disease have fallen 40 percent from 1999 to 2009 and have dropped for stroke nearly 37 percent during that same time period.[1] This decline is directly related to heart and stroke research, with scientists on the verge of new and exciting discoveries that could lead to innovative treatments and even cures for

Attention should also be given to other 20 to 22 NIH institutes (out of 27), centers and divisions that conduct heart and stroke research, primarily the:

Å National Institute on Aging (NIA): the NIA leads the federal effort supporting and conducting research on aging and the medical, social and behavioral issues of older people;

Å National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK): the NIDDK conducts and supports basic and clinical research and research training on some of the most common, severe and disabling conditions affecting Americans. The Instituteôs research interests include: diabetes and other endocrine and metabolic diseases; digestive diseases, nutrition, and obesity; and kidney, urologic and hematologic diseases; and

Å National Institute of Nursing Research (NINR): the NINR supports basic and clinical research that develops the knowledge to build the scientific foundation for clinical practice, prevent disease and disability, manage and eliminate symptoms caused by illness, and enhance end-of-life and palliative care.

physical education courses to meet the needs of disabled students.
Do not allow students to opt out of physical education to prepare for other classes or standardized tests.
Schools should implement programs to support evidenced-based physical education, activity, and fitness,

communities, especially in predominantly Latino or African-American neighborhoods, often have fewer resources to support active lifestyles and places to play and exercise.[54] Programs targeted to low-income, racially and ethnically diverse populations can increase active commuting and are associated with higher overall levels of moderate to vigorous physical activity throughout the day.[55] Community-based physical activity interventions are cost-effective, reducing new cases of many chronic diseases and improving quality of life.[56]

There are various ways to promote physical activity and active transportation in the community:

Street Level Design/Complete Streets policies consider the needs of all users in all transportation projects incorporating walking, bicycling, public transportation, and driving.

Smart Growth Design communities are designed with active living as the focus. Communities are connected with street patterns that make it easy to walk or bike to destinations. Developers try to locate essential services like schools and stores closer to homes to encourage walking and provide green spaces for recreation.

Shared Use of School Facilities

Promote physical activity through shared use of School recreational spaces
Support physical activity opportunities through economic and other tax incentives such as tax relief for development of

Research documents the link between more frequent eating out and increased body weight.[66,67] Obesity is not only a

American Heart Association Recommendations for the Dietary Guidelines for Americans
The American Heart Association will actively engage throughout the entire process of updating and revising the Dietary Guidelines for Americans to assure that the best possible evidence base is informing the final policy document and providing guidance to improve the cardiovascular health of the US population.

Reducing the Marketing and Advertising of Unhealthy Foods to Children

Inappropriate consumption of low nutrient, high calorie foods contributes to energy imbalance and poor health. Additionally, electronic media use is significantly correlated with childhood obesity[85,86,87,88] and advertising unhealthy foods contributes to childrenôs food preferences, requests, and diet.[89,90,91,92] Even children up to the age of 12 have a difficult time identifying the persuasive intent of food advertising and marketing.[93] Consequently, the American Heart Association sees no health, ethical, political, scientific, or social justification for marketing and advertising low-nutrient, high-calorie foods to children and suppo

American Heart Association Policy Recommendations for Reducing Unhealthy Food Marketing and Advertising to Children

Ultimately advocate for federal regulatory oversight of foods marketed and advertised to children since voluntary standards have not shown a significant impact on health outcomes.[102][103]

In the interim, work to strengthen voluntary initiatives to include more participation by industry, a more comprehensive definition of the media/technology that is used to target children, and robust enforcement of the nutrition standards.

There should be no unhealthy food and beverage advertising to children in schools, on buses or education materials because children should have a learning environment free of commercial influence and pressure.

Supports measures that restrict food advertising and marketing to children including, but not limited to allowing only healthy foods to be marketed and advertised to children, discouraging the product placement of food brands in multiple media technologies, eliminating the use of toys in unhealthy kidsô restaurant meals, and using licensed characters on only healthy foods.

intakes of milk, calcium, and other nutrients and an increased risk of several medical problems including diabetes.[133,135,136,137]

Maintain funding for SNAP Nutrition Education and strengthen the program to ensure low-income Americans have the ability to make healthy choices, increase fruit and vegetable consumption, and reduce their risk of chronic disease and obesity. Incorporate physical activity into SNAP education

Establish a multistate incentive pilot to promote increased SNAP purchases of fruits and vegetables at farmers markets and other healthy food retailers and support private/public partnerships and programs like Wholesome Wave, Philly Food Bucks, Roots of Change and Double Up Food Bucks that have shown great success[180]

Maintain funding for and promote the Agricultural Marketing Service's Farmers' Market Promotion Program (FMPP). The FMPP is a competitive grant program that makes funds available to eligible entities for projects to establish, expand, and promote farmers markets, roadside stands, community-supported agriculture programs, agritourism activities, and other direct producer-to-consumer opportunities

Provide grants and loans for value-added agriculture to develop the small and mid-sized processing and distribution systems needed to get products from family farmers into local, regional, and national markets

Increase the availability of fruits and vegetables in school meals and remove barriers which prevent local farmers from selling products to local schools.

Foster community-led approaches to improve consumer access to healthy and fresh foods in low income neighborhoods

Assure the USDA commodity program continues to increase the healthy foods that are provided to states and government feeding programs

Ensure the affordability of healthy and fresh foods for low-income families and seniors through purchases of fresh foods directly from farmers and other agricultural producers.

Provide incentives and crop insurance to small and mid-size farms to produce specialty crops like fruits and vegetables and distribute locally and regionally

Limit use of SNAP benefits in fast-food restaurants except for homeless, disabled, and some seniors

Support pilot programs with robust evaluation components to provide incentives to SNAP beneficiaries to purchase healthier foods

Support continued evaluation of all programs that provide local access to healthy, affordable foods

Recent advances include, but are not limited to, PDAs, handheld videogame consoles, mobile phones, smartphones, smartbooks, and portable media players. Their potential uses span from cellular conversation via voice, text, and visual media, to Internet access and digital software applications. Technological advances and improved computer processing power mean that single mobile devices such as smart phones and PDAs are increasingly capable of high level performance in many or all of these functions.

The functions of mobile health technologies that are apt for healthcare self-management are promoted by their useroted by

employee health in an environment where adults spend a large part of their time.[222] The Association also maintains that the use of rewards and penalties tied to health status should not jeopardize an employeeôs access to affordable, quality health care or be used as subterfuge for discrimination based on health status.

On May 29, 2013, the U.S. Departments of Health and Human Services, Labor and the Treasury issued final rules on employment-based wellness programs. The final rules incorporated feedback from numerous consumer groups and employers and support workplace health promotion and prevention as a means to reduce the burden of chronic illness, improve health, and limit growth of health care costs. Significantly, the final rules add additional consumer protections to ensure that these programs are not a form of medical underwriting where health costs are shifted from healthier employees to less healthy employees or where individuals are penalized if they have a preexisting condition or are genetically predisposed to a disease or risk factor. These consumer protections require that health-contingent wellness programs be reasonably designed, be uniformly available to all similarly situated individuals, and accommodate recommendations made at any time by an individualôs physician based on medical appropriateness.

The final rules continue to support ñparticipatory wellness programs,ò which generally are offered to employees without regard to an individualôs health status. These include programs that reimburse for the cost of membership in a fitness center, that provide a reward to employees for attending a monthly, no-cost health education seminar, or that reward employees who complete a health risk assessment, without requiring them to take further action.

The rules also outline standards for nondiscriminatory ñhealth-contingent wellness programs,ò which generally reward individuals who meet a specific standard related to their health. Examples of health-contingent wellness programs include programs that provide a reward to those who do not use, or reduce their use of tobacco, or programs that reward those who achieve a specified health-related goal such as cholesterol level, weight, or body mass index, as well as those who fail to meet such goals but take certain other healthy actions.

The final rules do allow significant flexibility for employers to design their own programs and allow employers to vary health care premiums/deductibles by 30% for achieving a health factor and up to 50% for tobacco use. The final rules will be effective for plan years beginning on or after Jan. 1, 2014.

A recent study by RAND, commissioned by the Department of Health and Human Services and the Department of Labor and authorized by the Public Health Service Act summarized a comprehensive review of the scientific and trade literature as well as a national survey, statistical analyses of health plan claims and wellness program data from several employers, and case studies of five employers with established wellness programs.[223] Key findings of the report were:

- Wellness programs are popular
- Program availability increases with employer size where larger employers have more extensive offerings (disease management, lifestyle interventions, more comprehensive screening).
- Typically, programs are implemented with some kind of screening combined with lifestyle intervention or disease management
- Employers find it fairly easy to get employees to participate in screening, but harder to participate in interventions. Program uptake is helped with healthy worksite culture, leadership role modeling and buy-in, and multiple communication channels.
- Employers are optimistic about the impact of wellness programs, however very few were able to provide cost and health impact data since

Health care costs level out for participants, but go up linearly in non-participants.
Financial incentives commonly used are in the $100 range. Employers are not anywhere near the currently allowed 20% variation when these incentives are tied to a health plan, the average is ~9%.
There is a small effect of incentives in promoting Health Risk Assessment completion (about 4% increase for every $25 invested). Rarely are incentives tied to disease management programs; smoking is the area where there are higher incentives for results (i.e. bigger sticks/and carrots). For other health factors, the difference between incentives for participation vs. outcome-based are not significant.
We cannot at this point conclude what effect incentives have on program participation, health outcomes, access to health care, and unintended consequences.
The database does not have more recent data (during the time that more outcomes-based incentives were implemented) and is not longitudinal enough to draw significant conclusions around incentives.

Healthy Early Child Care

The American Heart Association advocates for strong health promotion and obesity prevention programs in early childhood programs. Child care settings are an important environment for forming good health habits around children's dietary intake, physical activity, and energy balance and thus combating the childhood obesity epidemic.[224] The 2005 National Household Education Survey reports that 74% of all US children aged 3 to 6 years not yet in kindergarten were in some form of non-parental care, and 57% were in a center-based child care program making this an ideal setting for obesity-prevention interventions targeting this age group.[225] Furthermore, it has been reported that many children from low-income backgrounds consume 50%

Food advertising and marketing is another important causative factor in the obesity epidemic.[238] Exposure to food advertisements and industry marketing strategies produces substantial and significant increases in energy intake in all children and the rise is largest in obese children.[239] Aggressive advertising of high-calorie, low nutrient-dense foods contributes to higher consumption of those foods and should not be allowed in child care settings.

Preventing and controlling childhood obesity will require multifaceted and community-wide programs and policies with parents playing a critical role. One of the most important factors influencing childrenôs health behaviors are parentôs eating and physical activity behaviors and their level of education.[240,241] Parents are important role models and are largely responsible for physical activity opportunities, the type of food presented to young children, the portion sizes offered, and the emotional context in which food is eaten.[242] Successful intervention efforts must work directly with parents from the earliest stages of child development to support healthful practices both inside and outside the home.[243]

The American Heart Association and Nemours have launched Healthy Way to Grow, a technical assistance program for child care centers across the country aimed at decreasing obesity among children ages birth to five years old with inaugural funding provided by The William G. McGowan Charitable Fund. The program provides direct, hands-on assistance, customized training, resources and tools to support healthy lifestyles in child care environments. Components of the program include: developing and adopting a center wellness policy, providing training and technical assistance, engaging parents, and encouraging and recognizing progress towards best practices and policies for physical activity, screen time, food and beverage choices, and infant feeding. The American Heart Association will be able to support this programmatic and technical assistance effort with policy work around professional development for teachers and staff, credentialing and licensing, and nutrition, physical activity, and screen time standards for the early childcare environment.

Local Wellness Policies

The Child Nutrition and WIC Reauthorization Act of 2004 (P.L. 108-265, Section 204) required school districts participating in the National School Lunch Program (NSLP; [42 U.S.C.1751 et seq.]) or other child nutrition programs (42 U.S.C. 1771 et seq.) to adopt and implement a wellness policy starting with the 2006-07 school year. The Healthy, Hunger-Free Kids Act of 2010 (P.L. 111-296) extended this requirement and requires the U.S. Department of Agriculture (USDA) to develop regulations that provide a framework and guidelines for local wellness policies that include, at a minimum:

Million Hearts

Tobacco

Cigarette smoking continues to be the leading cause of preventable disease and death in the United States claiming approximately 467,000 lives prematurely every year.[1] Smoking not only claims the lives of those who use tobacco, but also those who are exposed to second-hand smoke. Smoking costs the U.S. economy more than $301 billion per year, including workplace productivity losses of $67.5 billion, premature death at $117 billion, and direct medical expenditures of $116 billion.[252] Tobacco control efforts by the American Heart Association have contributed to a decline in U.S. cigarette consumption by more than 24% over the last decade. [253] Despite this progress, 21.3 percent of men and 16.7 percent of w1e.

estimated at $10 billion each year.[265] If recent trends in the reduction in the prevalence of passive smoking continue, the health and economic burden of passive smoking in the U.S. would be cut annually by approximately 25%ï30%.[9] This potential reduction has important ramifications for lowering Medicare, Medicaid, and private insurance costs.

There are other important economic arguments. The hospitality and tobacco industries often promote the idea that business will suffer after these laws are passed. However, increasing evidence from

portion of the housing costs.

Surveillance data show that the smoking rate is higher in subsidized housing where 32.7% of adults use tobacco compared with 20.6% in the general population.[254,274] As more states and localities have passed smoke-free air laws for public spaces and workplaces, the home is the most significant source of exposure to second-hand smoke, especially for children.[275] Americans on average, spend about two thirds of their time each day in their residences.[276] However, only half of U.S. households with both children and smokers have complete home smoking bans and unfortunately bans are less common among smoking families with older children, in African-American and Hispanic households, and in households in states where there is a higher smoking prevalence.[277]

Even if people living in multi-unit housing have a smoke-free policy for their own home, they may still suffer incursions from others in the complex. Research has documented the transfer of second-hand smoke in the air[278,279,280,281,282,283,284] and transfer of second-hand smoke constituents through heating, ventilation, air conditioning systems and other connections between units.[6,285,286,287] As many as half of multi-unit housing residents report that smoke has entered their unit from elsewhere in the building or complex[288,289] and detectable levels of nicotine have been documented in multi-unit buildings where smoking is permitted.[290,291,292]

In 2009, the U.S. Department of Housing and Urban Development encouraged smoke-free policies in public housing to prevent the migration of second-hand smoke between housing units in an attempt to lower exposure especially among the most vulnerable tenants including the elderly, children, and people with chronic illnesses.[293] In public housing, children and adolescents are 39 percent of residents while older Americans comprise 15 percent of residents.[294] There is evidence that exposure to second-hand smoke disproportionately affects minorities,[295,296] women, and those in lower socioeconomic groups since a larger number of these individuals are residing in subsidized housing.

One recent study[297] estimated the annual cost-savings associated with smoke-free policies in multi-unit housing by calculating savings for second-hand smoke related health care costs, renovations of housing units that permit smoking, and smoking-attributed fires. Renovations or repairs include paint to cover smoke stains, cleaning of ducts, replacing stained window fixtures, and replacing carpets. The calculations from this study showed that prohibiting smoking in all U.S. subsidized housing could save approximately $521 million per year, including $341 million in second-hand smoke-related hea

Tobacco products are taxed in different ways and at different rates, which has created large disparities in the tax levied on similar products. Such disparities have created opportunities for manufacturers to make small modifications to products or their labeling so that they qualify for lower tax rates ï including a recently revelation that cigar companies use kitty litter in their products to make them heavier and avoid higher taxes for lower-weighted products. The availability of these lower-taxed products can dissuade tobacco users from quitting and encourage youth to initiate tobacco use.

An April 2012 GAO report highlights how certain manufacturers have avoided paying higher taxes on roll-your-own tobacco by re-labeling the product as ñpipeò tobacco, which is taxed at substantially lower rates under the current tax code. The GAO also noted that some manufacturers have avoided the higher tax rates for cigarettes and small cigars by slightly modifying their products to qualify as large cigars. GAO estimates

Master Settlement Agreement, there could have been millions of fewer smokers just over a decade later.[129]

American Heart Association Priorities for Tobacco Prevention and Cessation Programs
The American Heart Association advocates for sustainable funding for state tobacco prevention and cessation programs to levels that meet or exceed CDC recommendations. Tobacco control programs should be comprehensive in accordance with CDC recommendations, staffed appropriately, and administered effectively with periodic evaluation Protect the Prevention and Public Health Fund

Advocate for Comprehensive Coverage of Tobacco Cessation Services in Private and Public Health Insurance

The American Heart Association advocates for comprehensive coverage of tobacco cessation services in public and private health insurance programs that includes medications and counseling. In general, tobacco cessation treatment remains highly cost-effective, even though it is very difficult for people to quit this deadly, addictive habit.[316] There is a strong relationship between the length of time patients have in behavior counseling sessions, the amount of time they are able to spend with their health care providers and successful treatment outcomes.

Advocate for strong regulations from the FDA to regulate all tobacco products, including cigars. Oppose any legislation that would promote or promulgate loopholes and exemptions for any tobacco products, including cigars.

The FDA has announced that it intends to assert authority over all tobacco products, including cigars. No tobacco product was excluded from FDA jurisdiction under the Family Smoking Prevention and Tobacco Control Act so that FDA could evaluate the science and public health considerations of every product. However, FDA has been slow to issue this regulation and as a result, tobacco companies are taking advantage to modify products to avoid regulation. In addition, legislation has been introduced in the past few Congress that would exempt certain types of cigars.

Consumption of cigars is rising. Sales of cigars more than doubled between 2000 and 2012 from six billion cigars to more than 13 billion cigars. Cigar consumption has been increasing while cigarette consumption has declined, and much of the growth can be attributed to smaller cigars that resemble cigarettes. While cigar smoking conjures images of middle-age and older men, todayôs cigar smoker is more likely to be a youth or young adult, and that number is growing. Results from the 2011 Youth Risk Behavior Survey show that more than one in six (17.8 percent) high school boys currently smoke cigars.[20] Each day, more than 3,000 kids under 18 years old try cigar smoking for the first time.[327]

Disparities in regulation have created opportunities for manufacturers to make small modifications to products or their labeling so that they qualify for lower tax rates ï including a recently revelation that cigar companies use kitty litter in their products to make them heavier and avoid higher taxes and regulation for lower-weighted products. They have also allowed companies to produce candy and fruit-

American Heart Association Priority for Eliminating Tobacco Sales in Pharmacies
The American Heart Association supports policies that prohibit the sale of tobacco products in all pharmacies

Air Pollution

Air pollution is associated with a variety of negative health outcomes, including increased risk of cardiovascular disease and stroke. Pollution is comprised of a mixture of substances from sources such as vehicle and power plant emissions and the burning of fossil fuels. The Clean Air Act requires the Environmental Protection Agency (EPA) to set National Ambient Air Quality Standards (NAAQS) for six ñcriteriaò pollutants: carbon monoxide, lead, nitrogen dioxide, ground-level ozone, particulate matter, and

Studies examining particulate matter and cardiovascular disease have found associations between an increase in $PM_{2.5}$ and out

those found in indoor air pollution.
Monitor opportunities to influence legislation and regulation at the state and federal level to decrease the amount of particulate matter air pollution from various sources.
Offer incentives to consumers and businesses to purchase energy-efficient technologies in order to reduce emissions from residential and commercial heaters, boilers, lighting, chillers, air conditioners, etc.
Promote ñgreen powerò markets, which allow consumers to purchase electricity generated by renewable sources.
Redesign utility rate structures to incorporate incentives for energy efficiency and clean energy.
Set building codes for new commercial and residential construction at a minimum level of energy efficiency. Specify requirements for ñthermal resistanceò in the building shell and windows, minimum air leakage, and minimum heating and cooling equipment efficiencies.
Incentivize power plants to install modern emission control systems; establish mandatory state or regional cap and trade programs which control power plant pollution by providing economic incentives for achieving reductions in the emissions of pollutants.
Invest in clean energy development, including wind and solar energy.
Set tougher motor vehicle standards for tailpipe emissions, enforce vehicle inspection and maintenance programs, and establish programs to help consumers purchase energy-efficient cars.
Establish a diesel emissions reduction fund, aggressively retrofit and replace state vehicles and equipment, adopt and enforce anti-idling ordinances and legislation, mandate closed crankcase ventilation systems, and promote truck stop electrification programs.
Promote alternative fuels for vehicles and equipment (including natural gas, propane, methanol, ethanol, electricity and biodiesel fuel) and regulate gas stations to reduce emissions.
Establish tolls, fuel fees, carpool lanes, and other programs that reduce the number of vehicle miles travelled on roads and increase funding for public transportation.
States can ñlead by exampleò by developing state energy plans, mandating renewable and energy-efficient purchase commitments for state facilities, offering loan and incentive programs to increase energy efficiency in public buildings, retrofit and replace state vehicles and equipment, and implement a public communication strategy regarding the benefits of clean energy.

IV. Access to Appropriate and Affordable Health Care

Expand and protect access to affordable, adequate, transparent insurance coverage for a*ll*

Implement and Build on the Affordable Care Act

The associationôs work to implement the coverage provisions of the Affordable Care Act continues to be guided by our first principle for healthcare reform, that all residents of the United States should have meaningful, affordable healthcare coverage. The Congressional Budget Office projects that 30 million Americans will gain access to coverage once the law is fully implemented, making the ACA the most important piece of legislation for expanding access to care since Medicare and Medicaid were created. Implementation of the law will need to continue to be monitored closely and the association will need to continue working to ensure that insurance coverage is indeed

expansion in states that have not yet done so, addressing barriers to coverage for undocumented residents, and eliminating or reducing tobacco surcharges that may make coverage prohibitively expensive for tobacco users.

Protect Medicare, Medicaid, and CHIP

Medicare, Medicaid, and the Childrenôs Health Insurance Program are important sources of insurance coverage for adults and children with or at-risk for CVD and stroke. Medicaid alone provides an important safety net for 16 million Americans with a history of heart disease, stroke or other forms of cardiovascular disease (CVD), including seniors living in nursing homes, children with congenital heart disease, and those who have been disabled by stroke, congestive heart failure or other CVD. In addition, 42 percent of Medicare beneficiaries have a heart condition and 12 percent have had a stroke. As the federal and state governments struggle with budget deficits, Medicare and Medicaid in particularly are coming under increasing attention as a potential source for budget savings, even as the Baby Boomers age and more people need the coverage these programs provide. As Congress considers changes to these public programs, the association will work to:

Protect access to Medicaid and Medicare for the millions of Americans with heart disease and stroke.

Maintain the long-term sustainability of these two programs which are essential to our patient populations.

Make the impact on patients, particularly those who are most vulnerable, the central focus of any dialogue on health entitlement reform.

Ensure that entitlement reforms emphasize improvements in health care value, rather than shifting costs from the public to the private sector or from the government to beneficiaries.

Support changes that promote prevention and coordinated care and reward higher quality as the best approaches to achieve significant cost savings and improvements in health outcomes.

Access to Stroke Rehabilitation

The association works to protect and improve stroke survivorsô access to rehabilitative services in Medicare, Medicaid, and private insurance coverage. Examples of the types of policies we continue to support include:

Actively advocating for Congress to repeal the Medicare outpatient therapy caps for physical and speech therapy and for occupational therapy or to extend the ñexceptions processò that Congress has put in place to allow Medicare beneficiaries who need medically necessary therapy services to get an exception from the caps.

Advocated for the inclusion of ñrehabilitative and habilitative services and devicesò as one of the 10 categories of ñEssential Health Benefitsò that have to be covered by all private health plans in the nongroup and small group markets and for many Medicaid beneficiaries, starting January 1, 2014. As this coverage is implemented, the association will need to continue monitoring it to ensure that access to therapy services is sufficient and that limits on therapy services are not preventing access to needed rehabilitative and habilitative care.

Increase Access to Cardiac Rehabilitation

Each year, roughly 785,000 Americans will have a heart attack and more than 60 percent will have a second and potentially fatal event.[1] Cardiac rehabilitation (CR) reduces the risk of a future cardiac event by stabilizing, slowing or even reversing the progression of cardio-vascular disease (CVD).[379] Patients with other cardiovascular diseases such as valve repair and heart failure also benefit from exercise rehabilitation.

Yet despite its clear benefits, CR remains underutilized, particularly among women and minorities.[380,381] Only 14% to 35% of eligible heart attack survivors and 31% of patients after coronary bypass surgery participate in a CR program.[3,4] The utilization rate for eligible Medicare beneficiaries is an even lower 12%, and evidence clearly

T

Delivery System Transformation: Medical Homes & Accountable Care

The Patient Centered Medical Home (PCMH) model is emerging as a key vehicle to transforming the organization and delivery of primary care, improving health care quality, and controlling costs.[421,422] The PCMH is a whole-person orientation to patient care that is responsible for meeting the large majority of patientsô physical and mental needs, including prevention, wellness, acute care and chronic care.[423,424] PCMHs emphasize integrated and coordinated care, focus on quality and safety, facilitate partnerships between patients and physicians, and empower patients with strategies for self-management.[423,424,425]

Evaluations of PCMHs in the private sector reveal that these models of care are meeting cost, utilization, and quality objectives by reducing hospital admissions and use of emergency department services, lowering medical and pharmacy costs, and achieving more efficient care delivery.[426,427,428,429,430] One recent evaluation demonstrated up to $4.5 return on investment for every $1 spent on developing a PCMH.[427]

The ACA calls for the broader application of shared decision-making as a way to improve quality and patient experience.[464,465,466]

Value-Based Insurance Design

Value-based insurance design (VBID) is another tool that has emerged to reduce health care spending. VBID aims to increase health care quality and decrease costs by using financial incentives (such as reducing or eliminating co-pays) that encourage consumers to select high-quality, cost-effective health care services ï those services that yield health benefits of high value relative to their costs. [467,468] By designing insurance packages that offer preventive care, wellness visits and certain high-value treatments (such as medications to control blood pressure) at little or no cost to consumers, health plans can promote prevention, healthy behaviors and treatment adherence among beneficiaries, all which may save money by reducing future expensive medical procedures.[469,470] Benefit plans may create disincentives as well, such as high cost-sharing, for health choices that may be unnecessary or repetitive, or when the same outcome can be achieved at a lower cost.

ACA emphasizes elements of VBID. For example, the ACA now requires private insurers and medical expansion plans to cover certain clinical preventive services without cost-sharing,[471,472] many of which are important for CVD prevention.[473] With careful design, implementation, and evaluation, value-based cost sharing can be an important tool for aligning patient and provider incentives to pursue high-value care. Recent research has demonstrated that reducing cost-sharing for certain drugs and treatments (such as reduced co-pays for statins and beta-blockers) has the potential to yield improved care and reduced costs for patients with CVD.

Innovation Center. These initiatives will help, but greater investment is needed. For example, the Association of American Medical Colleges and the Association of Academic Health Centers, among others, have advocated for changes in medical school curricula to help better train cardiologists and other specialists to work with non-physician professionals (e.g. nurses, family caregivers etc.).[509,510,511,]

Nursing: Scope of Practice and Training. In many states, scope-of-practice laws limit nurse practitionersô (NP) capacity to practice to the fullest extent of their education, training, and competence.[400,487,512] For example, many states do not allow NPs to see patients, prescribe medications, or order and evaluate tests without physician supervision.[513] These laws ï which often are not based on education-level or safety concerns[513] ï set further constraints on an already stressed healthcare workforce system.

Several studies have demonstrated that NPs are capable of providing primary care services of equal or higher quality, and at lower costs as compared to physicians performing similar services.[400,501,514] There is also evidence that patients have comparable outcomes when care is administered by an NP.[515 ,516,517] As the delivery of health care for patients with chronic disease increasingly moves toward team-based, coordinated care models, this increases the need for nurses to be trained and empowered to meet new care challenges. Expanding and standardizing nursing scope-of-practice laws could promote more efficient and effective team-based care.[506,518,519] In addition, to meet growing primary care demands, nurses should be encouraged to obtain more advanced degrees and CVD and team-based leadership skill training.[507,520, 521]

Shortages of Healthcare Professionals in Rural and Underserved Regions. Shortfalls in the healthcare workforce affect everyone, but the impact is most severe for vulnerable and underserved populations. Approximately 20 percent of Americans live in rural or inner-city locations designated as health professional shortage areas (HPSA).[484,522] HPSAs are caused in large part by the tendency of practitioners to settle in regions with an already high concentration of physicians and other healthcare professionals.[523] Rural and underserved areas ï areas where patients have an increased risk of CVD and stroke[524,525] ï are subsequently left without a sufficient supply of

providers to underserved crea buamore poley ini tat-4(a)-13(n)6(d)-5( )-2(in)5(ce)-3(n)7(i)-iv(th)7(e)-s wb

sch a7(e)-18n18a)6(e)--8s)3)-2(d)-5(escr)-8(b)-5(ed)-7( )-b(co)-7(o)-5(y),eir

Therefore, the AHA supports the workforce with the goal of ensuring and increasing access to quality care:

- Provide sufficient public health and medical education funding and clinical training resources to improve chronic disease management, care coordination and patient-centered care support and promote the development of new models of care delivery, including those that emphasize team-based approaches using allied health professionals.
- Monitor and pursue opportunities at the state level to address workforce capacity and access.
- Support the workforce needed for both healthcare delivery and research
- Promote the growth and diversity of the healthcare workforce through a sustained and substantial national commitment to medical education and clinical training
- Support healthcare reform proposals that provide sufficient public health funding, medical education funding and clinical training resources for programs that improve chronic disease management, care coordination and patient-centered care.
- Support healthcare reform efforts that promote the development of new models of care delivery, including those that emphasize the roles of allied health professionals.
- Support efforts to produce an adequate number of well-trained cardiologists who will devote themselves to prevention, early and accurate diagnosis, and cost-effective treatment of cardiovascular diseases.
- Promote efforts to produce advanced practice providers with specific training and experience in cardiac conditions that will contribute to cost-effective care, improved staffing, and continuity of care.

Given the prominence of USPSTF recommendations for coverage, it is important for AHA and other stakeholders to have a voice in the USPSTF recommendation process. Review and comment by stakeholders outside the Task Force occurs at several different points during the development of evidence reports and recommendation statements: (i) during development of a research plan, USPSTF incorporates expert review and partner organization[548] comment on background documents to confirm that all relevant outcomes are being considered, that relevant literature has been considered, and that the evidence presented for USPSTF consideration is accurate;[548] (ii) once draft recommendations are developed, they are posted on the USPSTF website for public comment, and distributed to all federal and primary care partners of the USPSTF as well as appropriate clinical specialty societies for input;[548] and (iii) stakeholders can also nominate a topic for consideration[549] or request reconsideration (due to new evidence, new tests available, or status changes in the public health burden of the condition) through the Task Forceôs website.[550]

There are also informal opportunities for review, where stakeholders can seek meetings with individual USPSTF members, partner organizations or federal partners. For example, one of the Task Forceôs new duties established under the ACA is ñimproved integration with Federal Government health objectives.ò[551] Consistent with this duty, stakeholders may urge certain federal agencies to request meetings with the Task Force to discuss specific federal initiatives, such as the *Million Hearts Campaign,* and how USPSTFôs work could contribute to advancing federal goals.

Community Preventive Services Task Force

The ACA also codifies the Task Force on Community Preventive Services (ñCommunity Task Forceò), which presents its recommendations on community-based prevention and health promotion activities in *The Guide to Community Preventive Services* (The Community Guide).[552,553] The Community Task Force is an independent, nonfederal panel of public health and prevention experts that provides evidence-based findings and recommendations about community preventive services, programs, and policies to improve health. To develop its recommendations, Community Guide scientists lead or support interdisciplinary teams that carry out systematic literature reviews to identify what works to promote health and prevent disease, injury and disability.[554,555] Evidence compiled and recommendations developed by systematic review teams are brought to the full Community Task Force for their consideration and approval.

While there is no formal structure in place to allow for public comment or stakeholder input into the review process, systematic review teams frequently collaborate with CDC scientists and program managers, federal and non-federal liaison organizations,[556] and other researchers, practitioners, and policymakers.[554,557] AHA has collaborated with systematic review teams in the past, and, in circumstances where AHA is not a member of a given review team, it can informally collaborate with individual systematic review team members or liaison organizations to submit input.

It should be noted that while many of the Community Guide recommendations are important for CVD prevention (for example, recommendations on CVD prevention and control,[558] obesity prevention,[559] nutrition and physical activity,[560] and tobacco smoke[561]), recommendations from the Community Guide do not carry the same weight as USPSTF recommendations: there is no requirement that these recommendations become integrated in to private and public health insurance coverage. The Community Task Force operates independently of USPSTF, though ACA requires USPSTF to ñtake.

Support Comprehensive

Continue operation of the NEMSIS Technical Assistance Center (TAC) to expand the number of EMS response and patient outcome records housed in the national EMS database.
Improve linkage capabilities with other health databases and state trauma registries while integrating NEMSIS with electronic health records to enhance patient care outcomes.
The creation of regular reports providing national data for providers and policymakers.

Stroke and STEMI Systems of Care

Time is of the essence in treating acute cardiovascular conditions, but in far too many cases a fragmented and disorganized delivery system prohibits patients from receiving the treatments that can improve or even save their lives. The AHA believes that leadership and resources at the federal, state, and community levels are needed to help develop and implement coordinated systems of care for acute conditions to improve patient outcomes.

Certain care processes have been demonstrated to improve patient outcomes. Systems of care seek to implement these processes so that care is coordinated and victims of heart attack receive timely and appropriate treatment.

Although the clot-dissolving drug tPA is available to treat the most common type of stroke, only 3-8.5% of eligible stroke patients receive this therapy. This thrombolytic therapy can significantly reduce disability from stroke, but should be administered as soon as possible after symptom onset and preferably within four and a half hours to be most effective. It also saves money by improving patient outcomes and reducing the need for more extensive medical care. A recent analysis of more than 2,700 stroke patients in the U.S. and Europe confirmed that greater use of tPA could save nearly $50 million per year in the U.S. alone. Initial efforts at implementing stroke care systems have shown that they improve patient access to recommended care, including increased administration of thrombolytic therapy.

Roughly 1 out of every 4 heart attack victims each year will have the most severe type of heart attack called ST-Elevation Myocardial Infarction (STEMI). The quicker a patient with this heart attack has the completely blocked artery reopened (ñreperfusionò), the better the chances are for survival and less permanent damage to the heart. Approximately one-third of STEMI patients do not receive any reperfusion therapy (including the preferred type, percutaneous coronary intervention (PCI)) to restore blood flow in the artery. Many more do not receive this treatment within the recommended 90 minutes, even though such treatment greatly reduces the risk of death or debilitation

Systems of care are tailored by states or regions and locally implemented to meet the needs and challenges of an area, but should be based on the latest scientific guidelines. The ideal system of care provides patients with seamless transitions from each stage of care to the next (EMS; acute care; subacute care secondary prevention; and rehabilition and coverage). There are gaps and needs at each stage of this care continuum, however, that could be addressed by more coordinated care.

The American Heart Association/American Stroke Association advocates for resources and policies to help facilitate the development of coordinated systems of care in states and regions for stroke and STEMI. Key approaches include:

Through an integrated approach with other AHA/ASA program activities, promote efforts to create inclusive and coordinated statewide systems of care to improve the treatment of the stroke patient.
Work to ensure that the recognition, and the protection, of Primary Stroke Center, Comprehensive Stroke Center and Acute Stroke Capable Facility designation is based on Joint Commission/AHA/ASA accreditation or an equivalent accreditation process offered by a guidelines-based, nationally recognized accrediting organization.
Utilizing current AHA/ASA guidelines for stroke care, promote within EMSS statewide standardization and implementation of stroke training, assessment, treatment, and transportation protocols.
Support access and coverage of rehabilitation services for stroke patients.

Through an integrated approach with the Mission: Lifeline initiative, promote efforts to coordinate systems of care to improve the treatment of heart attack and sudden cardiac arrest patients by adhering to ACC/AHA guidelines and are consistent with Mission: Lifeline recommendations for criteria for heart attack and sudden cardiac arrest systems of care.

content and timeliness of communication from EMS to hospitals should be addressed to proactively mobilize healthcare personnel before arrival and reduce time delays to treatment;

Regional systems may involve a town, a city, a county, a state, or another region of the country. Systems should include academic or community receiving hospitals with multidisciplinary teams, including cardiology, critical care, and neurology. The volume of patients who have restoration of circulation after cardiac arrest is not solely tied to institutions but to practitioners who practice at multiple institutions;

Referral hospitals will continue to play a vital role in optimizing care for patients with restoration of circulation after OOHCA. Their immediate efforts, before transfer to the receiving hospital, in initiating therapeutic hypothermia early in conjunction with EMS will be important in the final outcomes of many patients. Referral hospitals should be provided with the necessary funds for equipment and education and be required to follow specific patient care and triage protocols, and they should report their experience, as has been done in selected inclusive regional trauma systems;

As with trauma centers, burn centers, STEMI centers, and stroke centers, national criteria should be developed to enable the categorization, verification, and designation of centers for the treatment of patients with restoration of circulation after OOHCA. External credentialing should be required as opposed to self-designation to support the development and sustainability of adequate patient volumes and high-quality care. The number of level 1 cardiac resuscitation centers in a given region should be limited to maintain provider skill levels and to justify the initial costs and institutional commitment required to care for these specialized patients;

Assessments of provider or hospital performance of acute coronary angiography should separate procedures performed in patients resuscitated from cardiac arrest from those performed in other patients to reduce potential disincentives to the performance of an intervention in these patients with high morbidity and mortality;

Evidence-based best practices and model EMS protocols should also be developed to guide states and local EMS systems in developing inclusive regionalized approaches to post-resuscitation care.

The American Heart Association/American Stroke Association advocates for public policies and resources that support a comp

Telehealth
Reimbursement
Ease licensing/credentialing barriers

V. Promote High Quality, High Value Health Care

In its landmark 2001 report, Crossing the Quality Chasm, the Institute of Medicine (IOM) declared that, ñBetween the health care we have and the care we could have, lays not just a gap, but a chasm.ò[570] Shortly thereafter, McGlynn and Asch estimated that Americans only receive the recommended care approximately half of the time.[571] The report and this subsequent quantification served as a call to action for the healthcare community to focus its attention on improving patientsô receipt of evidence-based care. Recent policy discussions and activities, spearheaded by the passage of healthcare reform, have focused on improving healthcare quality, as one component of what has come known as the ñtriple aimò - better health, better health care, and care at a lower cost. Together this tripod represents a high-value healthcare system with the aim of getting the highest quality and improvement in health out of every healthcare dollar spent.

This section provides an overview of the AHAôs public policy approach to improving healthcare quality, including that which improves adherence to clinical guidelines and care protocols, promotes safe, evidence-based diagnosis and treatment of cardiovascular disease and stroke, and reduces health inequities.

Adherence to Clinical Guidelines and Treatment Protocols

A ñclinical registryò is a database of health information on specific clinical conditions, procedures or populations. The data collected in a registry captures clinically important events relevant to a particular population or condition. Registries can be integrated with EHRs to directly support evaluation of care delivery and patient outcomes. Registries can broaden knowledge of clinical service patterns, processes and patient outcomes and capture valuable, real-time patient data that is not present in an administrative record, which typically only contains claims data or billing information. Because a registry can continuously capture data, registries have the potential to identify unnecessary or inappropriate variation and drive quality improvement by creating a continuous feedback loop to pinpoint areas of poor quality. In this way, the data collected and aggregated by the tool allows the provider or facility to identify problems with particular types of care, develop quality improvement interventions based on the identified problems, and monitor progress after implementation of a chosen intervention. As a result of the GWTG-Stroke program that uses a registry to collect and feedback information to participating hospitals, participating hospitals have shown improvement in adherence to stroke performance measures.[575]

Additionally, registries, as demonstrated by the experiences of the GWTG-Stroke registry can help hospitals reduce disparities in the care they deliver. While evidence suggests that not only are minorities at higher risk of suffering a stroke but they also receive lower quality of care and have worse health outcomes. Hospitals participating in GWTG-Stroke improved care for black, Hispanic, and white patients.[576] Clinical registries also support quality and safety evaluation by monitoring adverse events related to particular therapies, drugs, or devices[577,578] and examining provider adherence to safety protocols and best practice guidelines.[579]

Data collected via a registry can also catalyze systems changes on a regional or statewide basis to promote a more comprehensive and coordinated approach to care. Regional participation in quality improvement programs that utilize a clinical registry can help illuminate problems that exist in the system of care so that corrections and improvements can be made. For instance, data may show poor patient education about symptoms, geographical differences in the quality of care received, or problems with adherence to treatment guidelines. The data can then catalyze stakeholders to find solutions to the challenges encountered. For example, data from Marylandôs statewide stroke registry showed that some of the stateôs hospitals were reluctant to give tPA ï a drug used to treat thrombotic and embolic stroke ï to stroke patients. Having identified this concern, stakeholders were able to investigate its cause and determined that these hospitals were wary of prepping tPA because of the drugôs cost in the event that it was ultimately not used. Consequently, they developed a system where hospitals can return unused tPA to the manufacturer; this has promoted the delivery of tPA to appropriate stroke patients.[12]

The increasing sophistication of these tools and the pace of advances in health information technology offer significant promise for continued improvement in and better informed clinical decision-making.

American Heart Association Advocacy Priorities around Health Information Technology

Support public policies that encourage the development and implementation of health information tools, such as clinical-decision-making technology that deliver clinical guidelines in real-time to clinical decision-makers.

Promote policies that empower consumers to make informed decisions regarding the importance of owning, managing and maintaining personal health records.

Promote legislation and regulations that encourage the development and use of HIT with appropriate patient privacy safeguards.

Urge policy makers to create federal, state and local CVD and stroke registries in order to monitor incidence and support the development of relevant quality improvement initiatives.

Encourage policy makers to use patient-centered, evideQrc,ntinued-13(n)6(t)-10(s)3( )-2(r) ote ed, patient

Quality and Performance Measures

Quality measures attempt to represent the achievement of a recommended course of treatment by a healthcare practitioner by quantifying the existence of a structural component believed to connect to high quality care, a process understood to constitute high quality care, or quantify a particular set of outcomes. Performance measures are quality measures that are applied to the performance of a clinician or group of clinicians to either benchmark care delivery or measure improvements in quality over time. Quality-of-care measures can help create learning environments for health care professionals and ensure that best practices are applied uniformly to all patients.

ñPay for performance programsò are reporting programs that link the achievement of measure to payment whether as a financial reward through additional payment or a bonus or penalties via a reimbursement withhold. More research is needed to understand the impact of these programs on patient outcomes, both intended and unintended. Evaluation, ideally using clinical rather that administrative or claims data should be a central component of any program to better understand the programôs outcomes.[580] Given that many programs are still in their early stages of implementation which makes systematic evaluation and understanding their long-term impact difficult. A recent study, however, based on a cluster-randomized trial and published in the Journal of the American Medical Association did show an association between pay-for-performance programs and modest improvements in

CCHDs represent about 25% of all congenital heart defects.[586]

behalf of cardiovascular disease and stroke patients. In this way, it shows great promise in expanding the evidence base in these critical areas.

American Heart Association Priorities with Comparative Effectiveness Research
Promote the conduct and interpretation of comparative effectiveness research according to fundamental scientific principles.
Encourage funding for organizations, including PCORI, that support CER that aligns with Association principles.

Delivery System and Payment Reform

Delivery system reforms and a focus on testing structural and organizational models of care delivery, as well as their impact on care access, utilization and effectiveness have garnered recent focus. A recent Medicare study found that the average primary care practice coordinated with 99 other physicians working across 53 different practices.[588] In this way, many models focus on coordinating care among these providers, supporting medical documentation that makes health information portable, and disseminating information throughout the broader system to support proactive care management. Examples of specific models include the patient centered medical home (PCMH) and accountable care organizations (ACOs).

Payment models are often used in tandem with systemic or structural changes to create financial incentives for providers, patients, and others to encourage care delivery by clinicians and adherence by patients. These payment methods can include capitation, bundled payment, or care coordination/ patient management fees. Similarly to the work that it has done on pay-for-performance programs, the AHA/ASA believes that any payment reform program must have the goal of reducing the burden of disease for patients and developed a set of principles that aim to ensure that this end goal is met and unintended consequences do not occur.(should be numbered the same as January 2006 Bufalino citation, cited above) The principles stress the importance of transparency to consumers of the existence of a payment incentive to physicians and any resulting restrictions on access to services this incentive may create; the need to allow for variation so that the unique needs of sub-populations are met, as well as existing disparities in care delivery are addressed; and the use of quality measures that are evidence-based, meaningful, and designed to ensure accountability for quality, not just cost.[589] Additionally, any incentive used should encourage the development of enabling structures within the healthcare system that enhance its safety, effectiveness, efficiency, equity, timeliness, and patient-centeredness.[590]

Further testing and evaluation of these models is needed to understand whether they are effective, as well as any unintended consequences occur that need to be avoided. The Affordable Care Act created the Center for Medicare and Medicaid Innovation (CMMI) within the Centers for Medicare and Medicaid Services (CMS) to conduct this testing and evaluation and gave it the authority to scale these models without additional Congressional action if evaluation showed they improved quality while reducing or maintaining cost.[591]

American Heart Association Priorities for Delivery System and Payment Reform
Monitor existing demonstration models testing new delivery and payment reforms.
Encourage adoption of evidence-based methods of care coordination.
Encourage researchers to examine and evaluate the impact of care delivery models on cardiovascular and stroke treatment and patients, with particular attention to identifying unintended consequences as they occur.
Evaluate the role of quality in health care payment systems.
Support research and evaluation of mechanisms for aligning payment with healthcare improvement.

Benefit design

Health insurance benefit design is also increasingly being used to adjust both provider and patient behavior, making it important that public policy supports appropriate patient safeguards in the development and implementation of these designs, as well as balances the need for patient access to a wide variety of treatments with the affordability of available treatments.

The Association works with CMS, as well as other payors to evaluate the appropriateness of services and procedures, such as carotid artery stenting for stroke patients, anticoagulant home monitoring for patients with atrial

substitution, but opposes therapeutic substitution and supports communication of the therapeutic interchange to both provider and patient.

The Association supports policies that promote formularies that are consistent with the recommendations laid out in its position statement.

American Heart Association Priorities on Drug Formularies
Monitor, evaluate, and promote policies concerning drug formularies that are consistent with the AHAôs Drug Formulary position statement. Educate lawmakers about the need to balance access and cost considerations when developing formularies for public programs, such as Medicaid and SCHIP. Monitor drug formulary policy

The Association supports policies that promote formularies that permit therapeutic interchange and generic substitution when necessary and in designated circumstances and do not allow for therapeutic substitution. The AHA continues to monitor, evaluate and promote proposed public policies concerning drug formularies. The Association supports increased access to a broad range of cardiovascular disease and stroke drugs, treatments and medical devices and opposes therapeutic substitution of prescription drugs within a class. Public policy should support appropriate patient safeguards in the development and implementation of formularies. Further, drugs and devices used to treat or prevent cardiovascular disease and stroke must be properly reviewed, labeled, dispensed and marketed.

Medication Adherence

Nonadherence to prescribed medications is a growing problem that has become too common; the consequences include adverse health events and added costs to the U.S. healthcare system. The New England Healthcare Institute estimates that patients who fail to adhere to their medications cost the U.S. healthcare system $290 billion annually.[597] Patients with cardiovascular diseases contribute to poor adherence, raising the risk for heart disease and stroke. According to research by Vrijens and colleagues, roughly half of patients who have been prescribed high blood pressure medications stop taking them within a year.[598] Additional research shows that nearly one fourth of patients who have been hospitalized for an acute heart attack neglect to fill their prescriptions, only raising the risk of a subsequent heart attack.[599]

American Heart Association Advocacy Priorities for Drug Shortages
Support policies that foster communication between the FDA and manufacturing companies in order to prevent or mitigate drug shortages. Increase funding for research to better characterize cardiovascular care drug shortages and their impact on patient care. Educate and train healthcare providers on how to best mitigate shortages to avoid disruption in care.

Personalized Medicine

We are at the dawn of a new age, where our evolving knowledge of how genes and lifestyle combine to affect our health is transforming the practice of medicine. Genetic analysis can already identify risk factors for certain heart disorders, type 2 diabetes, and many other health conditions. Every few months, researchers are discovering more disease-related gene variants. This new information is being used to ópersonalizeô medicine according to each patientôs genetic profile. Genetic medicine can help identify disease risk as well as the likely response to certain drugs and drug doses. As scientists develop a greater understanding of the genetics of heart disease and stroke, we will move away from ñone-size-fits-allò medicine to more targeted and effective prevention, treatments, and even cures.

American Heart Association Advocacy Priorities for Personalized Medicine
Require disclosure of test validity and enact appropriate regulatory oversight for marketed tests to assure physicians and the public of test quality. Increase funding for research on the genetics of heart disease and the translation of new discoveries into preventive measures and treatments. Protect individuals undergoing genetic tests from discrimination by all forms of insurance. Educate and train the medical workforce to prepare for the expanded integration of genetics into healthcare practice. Integrate personalized healthcare with health information technology to deepen our understanding of the relationship between genetics, disease, treatments and outcomes. Ensure that intellectual property practices foster innovation and the development of new genetic tools. Educate the public about personalized medicine and encourage them to collect their family medical history.

Palliative Care

Palliative care is medical and supportive care for people with serious illness that is routinely integrated into care by all practitioners and focuses on providing patients and their families with relief from illness and suffering burden -- including symptoms, pain and stress--regardless of diagnosis[601]. By integrating medically appropriate care with supportive care practices, it helps patients and their families achieve goals of improved functioning and prolonging life, when possible, and with comfort and the preservation of hope at the end of life. Treatment goals are clarified and valued, care is coordinated across settings, and the patientôs practical, social, emotional, and spiritual needs are supported.

Research indicates that palliative care is beneficial for cardiovascular and stroke patients, given the symptom burden faced by these patients, the desire by many patients with serious illness (,)-2( )-2(ty)18(W*nBT3( )-2(l)-5(m)7(p)--2(s)3(er)-u)6(b) TJE

Research demonstrates that patients living with serious illness identify elements of palliative care such as pain and symptom management, avoidance of inappropriate prolongation of dying, achievement of a sense of control, and avoiding burdening others, as among their top priority needs from the healthcare system.[605] A majority of seriously ill patients, however, are not currently receiving palliative care.[603] As medical technology advances, patients are living longer and with conditions that were previously fatal, but with significant adverse implications for their quality of life and that of their families. Patients, who suffer from acute cardiovascular events or stroke when previously highly functional, also need additional support for coping and adjusting to the illness and complex decision making. Together, these factors support the need for the holistic approach taken by palliative care in

National Health and Nutrition Examination Survey (NHANES), Behavioral Risk Factor Surveillance System

The top areas of focus in the nonprofit sector are described below.

Support the Charitable Tax Deduction

More than 80 percent of the 46 million who itemized their tax returns in 2009 claimed the charitable deduction. These individuals are responsible for more than 76 percent of individual contributions to charitable organizations. Between 2003 and 2009, charitable organizations in the U.S. received $281 in online donations. More than 22 percent of those donations were made on December 30 and 31 each year; underscoring the extent to which donors are aware of, and influenced by, the tax implications of their giving. An April 2011 Gallup Poll found that 62 percent of Americans who do not claim the deduction support its preservation as an incentive for giving. While some believe the charitable deduction disproportionately benefits high income taxpayers, a 2012 study by the Center on Philanthropy at Indiana University found that 79.3 percent of high-net-worth households (annual income greater than $200,000) donated to basic needs charities in 2011.

Itôs important to remember that unlike incentives to save for retirement or to purchase a home, the charitable deduction encourages behavior for which a taxpayer receives no direct tangible benefit. It simply and effectively encourages taxpayers to give away a portion of their income to benefit others.

Encouraging Volunteerism

Public policy can play an important role in providing strong incentives for Americans to give their time in service to charitable organizations, including the American Heart Association. The AHA generally supports efforts to encourage people of

feeling of local ñownershipò of and engagement with the campaign, which in turn may lead to decreased participation and donations, especially for local charities.

2.

[104] Keller K. Forman J. Lee NM. Kuilema LG. Halford JC (2011). ñUse of licensed spokes-characters to increase intake of fruits and vegetables as part of a childhood obesity prevention program: pilot study results.ò Obesity. 19 Suppl. 1.
[105] Marino, C. J. and R. P. Mahan (2005). "Configural displays can improve nutrition-related decisions: An application of the proximity compatibility principle." Human Factors: The Journal of the Human Factors and Ergonomics Society 47(1): 121-130.
[106] Lewis, C. J. and E. A. Yetley (1992). "Focus group sessions on formats of nutrition labels." J Am Diet Assoc 92(1): 62-66.
[107] Geiger, C. J., et al. (1991). "Nutrition labels in bar graph format deemed most useful for consumer purchase decisions using adaptive conjoint analysis." J Am Diet Assoc 91(7): 800-807.
[108] van Kleef, E., et al. (2008). "Consumer preferences for front-of-pack calories labelling." Public Health Nutr 11(2): 203-213.
[109] Borgmeier, I. and J. Westenhoefer (2009). "Impact of different food label formats on healthiness evaluation and food choice of consumers: a randomized-controlled study." BMC Public Health 9: 184.
[110] van Herpen, E. and H. C. Trijp (2011). "Front-of-pack nutrition labels. Their effect on attention and choices when consumers have varying goals and time constraints." Appetite 57(1): 148-160.
[111] Hawley, K. L., et al. (2013). "The science on front-of-package food labels." Public Health Nutr 16(3): 430-439.
[112] Aschemann-Witzel, J., et al. (2013). "Effects of nutrition label format and product assortment on the healthfulness of food choice." Appetite 71C: 63-74.
[113] Grunert K (2007). ñA review of European research on consumer response to nutrition information on food labels.ò J Public Health 15, 385ï389.
[114] Wansink B (2003). ñHow do front and back package labels influence beliefs about health claims?ò J Consum Aff 37, 305ï316.
[115] Larsson, I., et al. (1999). "The 'Green Keyhole' revisited: nutritional knowledge may influence food selection." Eur J Clin Nutr 53(10): 776-780.
[116]Reid, R. D., et al. (2004). "The Heart and Stroke Foundation of Canada's Health Check food information program: modelling program effects on consumer behaviour and dietary practices." Can J Public Health 95(2): 146-150.
[117] Sacks, G., et al. (2009). "Impact of front-of-pack 'traffic-light' nutrition labelling on consumer food purchases in the UK." Health Promot Int 24(4): 344-352.
[118] Steenhuis, I., et al. (2004). "The effectiveness of nutrition education and labeling in Dutch supermarkets." Am J Health Promot 18(3): 221-224.
[119] Freedman, M. R. and R. Connors (2010). "Point-of-purchase nutrition information influences food-purchasing behaviors of college students: a pilot study." J Am Diet Assoc 110(8): 1222-1226.
[120] Liem, D. G., et al. (2012). "Health labelling can influence taste perception and use of table salt for reduced-sodium products." Public Health Nutr 15(12): 2340-2347.
[121] Sims, J., et al. (2011). Claiming health: Front-of-package labeling of childrenôs food. Prevention Institute; 2011.

[122] Vyth, E. L., et al. (2010). "Actual use of a front-of-pack nutrition logo in the supermarket: consumers' motives in food choice." Public Health Nutr 13(11): 1882-1889.
[123] Sutherland, L. A., et al. (2010). "Guiding stars: the effect of a nutrition navigation program on consumer purchases at the supermarket." Am J Clin Nutr 91(4): 1090S-1094S.
[124] Nestle, M. and D. S. Ludwig (2010). "Front-of-package food labels." JAMA: the journal of the American Medical Association 303(8): 771-772.
[125]

[131] Zenk, S. N. and L. M. Powell (2008). "US secondary schools and food outlets." Health Place 14(2): 336-346.
[132] Jha, P. and F. J. Chaloupka (2000). "The economics of global tobacco control." BMJ 321(7257): 358-361.
[133] Vartanian, L. R., et al. (2007). "Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis." Am J Public Health 97(4): 667-675.
[134]Flood, J. E., et al. (2006). "The effect of increased beverage portion size on energy intake at a meal." J Am Diet Assoc 106(12): 1984-1990; discussion 1990-1981.
[135] Nielsen, S. J. and B. M. Popkin (2004). "Changes in beverage intake between 1977 and 2001." Am J Prev Med 27(3): 205-210.
[136] Schulze, M. B., et al. (2004). "Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women." JAMA 292(8): 927-934.
[137] Nagai, Y., et al. (2009). "The role of peroxisome proliferator-activated receptor gamma coactivator-1 beta in the pathogenesis of fructose-induced insulin resistance." Cell Metab 9(3): 252-264.
[138] Jacobson, M. F. and K. D. Brownell (2000). "Small taxes on soft drinks and snack foods to promote health." Am J Public Health 90(6): 854-857.
[139] Mozaffarian, D., et al. (2012). "Population approaches to improve diet, physical activity, and smoking habits: a scientific statement from the American Heart Association." Circulation 126(12): 1514-1563.
[140]Storey, M. L., et al. (2006). "Beverage consumption in the US population." J Am Diet Assoc 106(12): 1992-2000.
[141] Levy, D. T., et al. (2004). "The effects of tobacco control policies on smoking rates: a tobacco control scorecard." J Public Health Manag Pract 10(4): 338-353.
[142] Andreyeva, T., et al. (2010). "The impact of food prices on consumption: a systematic review of research on the price elasticity of demand for food." Am J Public Health 100(2): 216-222.
[143] Duffey, K. J., et al. (2010). "Food price and diet and health outcomes: 20 years of the CARDIA Study." Arch Intern Med 170(5): 420-426.
[144] Block, J. P., et al. (2010). "Point-of-purchase price and education intervention to reduce consumption of sugary soft drinks." Am J Public Health 100(8): 1427-1433.
[145] Powell, L. M., et al. (2009). "Food prices and fruit and vegetable consumption among young American adults." Health Place 15(4): 1064-1070.
[146] Chaloupka, F. J. and L. M. Powell (2009). "Price, availability, and youth obesity: evidence from Bridging the Gap." Prev Chronic Dis 6(3): A93.
[147] Appel, L. J., et al. (2011). "The importance of population-wide sodium reduction as a means to prevent cardiovascular disease and stroke: a call to action from the American Heart Association." Circulation 123(10): 1138-1143.
[148] Whelton, P. K., et al. (2012). "Sodium, blood pressure, and cardiovascular disease: further evidence supporting the American Heart Association sodium reduction recommendations." Circula( )-2(Hea)-3(r)-5m[( )] TJE67ciation." a-os.ion ce 1c-2(cu

[212] Naughton, F., et al. (2012). "Randomized controlled trial evaluation of a tailored leaflet and SMS text message self-help intervention for pregnant smokers (MiQuit)." Nicotine Tob Res 14(5): 569-577.
[213] Soureti, A., et al. (2011). "Exploratory study of web-based planning and mobile text reminders in an overweight population." J Med Internet Res 13(4): e118.
[214] Blasco, A., et al. (2012). "Evaluation of a telemedicine service for the secondary prevention of coronary artery disease." J Cardiopulm Rehabil Prev 32(1): 25-31.
[215] Morikawa, N., et al. (2011). "Effect of salt reduction intervention program using an electronic salt sensor and cellular phone on blood pressure among hypertensive workers." Clin Exp Hypertens 33(4): 216-222.
[216] Merchant, R. M., et al. (2010). "Cell phone cardiopulmonary resuscitation: audio instructions when needed by lay rescuers: a randomized, controlled trial." Ann Emerg Med 55(6): 538-543 e531.
[217] Whittaker, R., et al. (2011). "A theory-based video messaging mobile phone intervention for smoking cessation: randomized controlled trial." J Med Internet Res 13(1): e10.
[218] Sakai, T., et al. (2011). "Effectiveness of the new 'Mobile AED Map' to find and retrieve an AED: A randomized

[237]Fitzgibbon, M. L., et al. (2005). "Two-year follow-up results for Hip-Hop to Health Jr.: a randomized controlled trial for overweight prevention in preschool minority children." J Pediatr 146(5): 618-625.
[238] Page, R. M. and A. Brewster (2007). "Emotional and rational product appeals in televised food advertisements for children: analysis of commercials shown on US broadcast networks." J Child Health Care 11(4): 323-340.
[239] Halford, J. C., et al. (2008). "Beyond-brand effect of television food advertisements on food choice in children: the effects of weight status." Public Health Nutr 11(9): 897-904.
[240] van der Horst, K., et al. (2007). "A systematic review of environmental correlates of obesity-related dietary behaviors in youth." Health Educ Res 22(2): 203-226.
[241] Krahnstoever Davison, K., et al. (2005). "Reexamining obesigenic families: parents' obesity-related behaviors predict girls' change in BMI." Obes Res 13(11): 1980-1990.
[242] Davis, M. M., et al. (2007). "Recommendations for prevention of childhood obesity." Pediatrics 120 Suppl 4: S229-253.
[243]Lindsay, A. C., et al. (2006). "The role of parents in preventing childhood obesity." Future Child 16(1): 169-186.
[244] Chriqui, J., et al. (2013). "School District Wellness Policies: Evaluating Progress and Potential for Improving Children's Health Five Years after the Federal Mandate. Brief Report. Volume 3." Robert Wood Johnson Foundation.
[245] Department of health and Human Services. Million Hearts. http://millionhearts.hhs.gov/index.html Accessed September 13, 2013.
[246] Sacco, R. L., et al. (2012). "What the million hearts initiative means for stroke: a presidential advisory from the American Heart Association/American Stroke Association." Stroke 43(3): 924-928.
[247] Tomaselli, G. F., et al. (2011). "The American Heart Association and the Million Hearts Initiative: a presidential advisory from the American Heart Association." Circulation 124(16): 1795-1799.
[248]Bikdeli, B. and J. A. Barreto-Filho (2012). "Reducing the cardiovascular disease burden: justified means for getting to the end." Circ Cardiovasc Qual Outcomes 5(4): 580-586.
[249] Afterschool Alliance. Afterschool Issue Overview. 2013; http://www.afterschoolalliance.org/researchFactSheets.cfm. Accessed April 19, 2013.
[250] Bassett DR. Erwin P. Fitzhugh EC. F

[291] Hood, N. E., et al. (2012). "Associations between self-reported in-home smoking behaviours and surface nicotine concentrations in multiunit subsidised housing." Tob Control.
[292] Matt, G. E., et al. (2011). "When smokers move out and non-smokers move in: residential thirdhand smoke pollution and exposure." Tob Control 20(1): e1.
[293]

[345]Xiang, H., et al. (2013). "Estimation of short-term effects of air pollution on stroke hospital admissions in Wuhan, China." PLoS One 8(4): e61168.
[346]

[397] Centers for Disease Control and Prevention. Heart Disease and Stroke Prevention: Addressing the Nationôs Killers. 2011. Available online: http://www.cdc.gov/chronicdisease/resources/publications/aag/pdf/2011/Heart-Disease-and-Stroke-AAG-2011.pdf
[398] Vincent, G. K. and V. A. Velkoff (2010). The next four decades: The older population in the United States: 2010 to 2050, US Department of Commerce, Economics and Statistics Administration, US Census Bureau.
[399] Leira, E. C., et al. (2013). "The growing shortage of vascular neurologists in the era of health reform: planning is brain!" Stroke 44(3): 822-827.
[400] Rodgers, G. P., et al. (2009). "ACC 2009 survey results and recommendations: Addressing the cardiology workforce crisis A report of the ACC board of trustees workforce task force." J Am Coll Cardiol 54(13): 1195-1208.
[401] Owens, D. K., et al. (2010). "AHRQ series paper 5: grading the strength of a body of evidence when comparing medical interventions ð Agency for Healthcare Research and Quality and the Effective Health-Care Program." J Clin Epidemiol 63(5): 513-523.
[402] Public Law 111-148, §6301.
[403] Alter, D. A., et al. (2009). "Relationship between cardiac rehabilitation and survival after acute cardiac hospitalization within a universal health care system." Eur J Cardiovasc Prev Rehabil 16(1): 102-113.
[404] Arrigo, I., et al. (2008). "Comparative outcome one year after formal cardiac rehabilitation: the effects of a random

[450] Whellan, D. J., et al. (2005). "Metaanalysis and review of heart failure disease management randomized controlled clinical trials." Am Heart J 149(4): 722-729.
[451] Bodenheimer, T., et al. (2002). "Improving primary care for patients with chronic illness: the chronic care model, Part 2." JAMA 288(15): 1909-1914.
[452] Chung, H., et al. (2013). "Early experience of a pilot intervention for patients with depression and chronic medical illness in an urban ACO." Gen Hosp Psychiatry 35(5): 468-471.
[453] Rokos, I. C. (2011). "Creating "turbo" accountable care organizations for time-critical diagnoses." Circ Cardiovasc Qual Outcomes 4(6): 647-649.
[454] Hibbard, J. H., et al. (2005). "Development and testing of a short form of the patient activation measure." Health Serv Res 40(6 Pt 1): 1918-1930.
[455] (2013). "Patient engagement, patient safety, and quality of care." Health Aff (Millwood) 32(2): 432-435.

[531] Aggarwal, B., et al. (2009). "Influence of caregiving on lifestyle and psychosocial risk factors among family members of patients hospitalized with cardiovascular disease." J Gen Intern Med 24(1): 93-98.
[532] Jaarsma, T., et al. (2009). "Palliative care in heart failure: a position statement from the palliative care workshop of the Heart Failure Association of the European Society of Cardiology." Eur J Heart Fail 11(5): 433-443.
[533] Mosca, L., et al. (2011). "Patterns of caregiving among patients hospitalized with cardiovascular disease." J Cardiovasc Nurs 26(4): 305-311.
[534] Piamjariyakul, U., et al. (2012). "Part I: heart failure home management: patients, multidisciplinary health care professionals and family caregivers' perspectives." Appl Nurs Res 25(4): 239-245.
[535] Piamjariyakul, U., et al. (2012). "Part 2: enhancing heart failure home management: integrated evidence for a new family caregiver educational plan." Appl Nurs Res 25(4): 246-250.
[536] Center for Healthcare Research and Transformation. Affordable Care Act Funding: An Analysis of Grant Programs under Health Care Reform. Sep 2012. Available at: http://www.chrt.org/assets/price-of-care/CHRT-Issue-Brief-September-2012.pdf.
[537] U.S. Preventive Services Task Force. About the USPSTF. Available at: http://www.uspreventiveservicestaskforce.org/about.htm.
[538] U.S. Preventive Services Task Force, ñUnderstanding How the U.S. Preventive Services Task Force Worksò (online at www.uspreventiveservicestaskforce.org/uspstf101_slides/uspstf101.htm).
[539] U.S. Preventive Services Task Force. Grade Definitions. February 2013. Available at: http://www.uspreventiveservicestaskforce.org/uspstf/grades.htm.
[540] U.S. Preventive Services Task Force. Recommendations. Available at: http://www.uspreventiveservicestaskforce.org/recommendations.htm.
[541] USPSTF A and B Recommendations. U.S. Preventive Services Task Force. Available at: http://www.uspreventiveservicestaskforce.org/uspstf/uspsabrecs.htm
[542] Patient Protection and Affordable Care Act, Public Law No. 111 at 148, § 1001, 124 Stat. 131, (March 2010), *amending* Part A of title XXVII of the Public Health Service Act, 42 U.S.C. 300gg et seq.
[543] Center for Medicaid and CHIP Services (2013). "Medicaid and Child43rd 3rdB Reco C1,

[558] Task Force on Community Preventive Services (2013). Community Preventive Services Task Force. Cardiovascular Disease Prevention and Control. Available at: http://www.thecommunityguide.org/cvd/index.html.
[559] Task Force on Community Preventive Services (2013). Community Preventive Services Task Force. Obesity Prevention and Control. Available at: http://www.thecommunityg

[581] Bardach, N. S., et al. (2013). "Effect of pay-for-performance incentives on quality of care in small practices with electronic health records: a randomized trial." JAMA 310(10): 1051-1059.
[582] Parker, S. E., et al. (2010). "Updated National Birth Prevalence estimates for selected birth defects in the United

[607] Chen, J., et al. (2011). "National and regional trends in heart failure hospitalization and mortality rates for Medicare beneficiaries, 1998-2008." JAMA 306(15): 1669-1678.
[608] Hauptman, P. J. and E. P. Havranek (2005). "Integrating palliative care into heart failure care." Arch Intern Med 165(4): 374-378.
[609]